RYAN CRENSHAW, M.D. Two-Day Preparation Instructions Prior to Colonoscopy Using SuFLAVE

Please read this packet, in its entirety, at least 2 weeks prior to your procedure. If you fail to follow the instructions and the procedure must be cancelled, the cancellation fee will be charged.

If you need to cancel your procedure, please let us know at least <u>5 business days</u> prior to the procedure. If you fail to do so, you will be charged a \$250.00 cancellation fee.

You will receive a confirmation email from our office staff at least 2 weeks prior to your scheduled procedure(s). All procedure(s) must be confirmed in order to remain on the schedule. If you do not receive a call from our office, please contact us at 703-444-4799.

If you take blood thinners such as Aspirin, Plavix, Xarelto or Coumadin, Dr. Crenshaw may recommend for you to hold these medications anywhere from 2-8 days prior to your procedure, depending on what agent you are taking.

Patients taking Aspirin: Please make sure one of the two lines is checked off below. If not, please contact our office at (703) 444-4799.

_	_ Please continue to take Aspirin, 81 mg or 325 mg, daily, including the day of the colonoscopy
	Please stop taking Aspirin, 81 mg or 325 mg, 8 days prior to the colonoscopy,

Any patient stopping Aspirin, Plavix, Coumadin, or any other blood thinner should contact the prescribing doctor (primary care physician, or cardiologist) to confirm that it is acceptable to stop this medication(s) for the recommended period of time. You may take Tylenol if needed. Please do not take any other medication or products that can thin the blood, such as Ibuprofen, Motrin, Advil, Aleve, Vitamin E and/or Garlic pills 8 days prior to the procedure(s).

It is highly recommended that you take your medication for heart disease, high blood pressure and asthma every day, including the day of your procedure. If you take any medication(s) around the time that you are taking a dose of the laxative to prepare for your colonoscopy, please take the laxative first, then your medication. When taking medication(s) on the day of your procedure, the medication(s) must be taken at least 4 hours prior to your procedure time with water. 4 hours prior to procedure time, you should stop consuming liquids, stop taking medications and not take anything orally until your procedure has been completed. All other medications should be brought to the hospital to be taken after your procedure.

If you are taking medications for diabetes, consult with the medical provider that is managing your diabetes to inform him/her that you are being asked to change your diet in preparation for colonoscopy. Please ask this provider how you should change your diabetes medication regimen to reduce the risk of your blood sugar becoming too low or too high during your preparation for the procedure.

You will need someone to drive you home from the hospital, or surgical center, after your procedure. You should not drive until the next day.

Two-day preparation instructions using SuFLAVE

(Polyethylene glycol 3350, 420 g, Sodium bicarbonate 5.72 g, Sodium Chloride 11.2 g and Potassium Chloride 1.4 g)

Caution: This preparation must be used with caution and may be contraindicated in patients with the following conditions: Kidney failure or compromise in kidney function, uncorrected electrolyte abnormalities, heart arrhythmias, history of seizure, impaired gag reflex, gastric retention, gastrointestinal obstruction and/or severe ulcerative colitis. If you have any of these conditions, please call the office to confirm this preparation is appropriate for you.

Five days prior to the procedure: Do NOT eat food containing seeds, corn, nuts, black pepper, lettuce, raw vegetables, fruits with seeds or skin as they can be difficult to lavage from the colon. Please do not take fish oil, krill oil, lemon oil or any supplements or foods that contain a significant amount of oil such as potato chips.

Two days prior to your scheduled colonoscopy, you may consume a normal breakfast. For lunch and dinner that evening, consume clear liquids from the menu of clear liquids listed below.

Please note that it is very important for you to be adequately hydrated during this process. There is no restriction on the volume of clear liquids that you consume. Consume as many of the clear liquids listed as possible to maintain adequate hydration and reduce the risk of electrolyte abnormalities during the preparation.

It is also important to vary the liquids you are consuming. Do NOT restrict yourself to drinking water only. Please make sure you consume liquids with salt such as soup broth. And liquids with carbohydrates/sugars such as apple juice. By varying the liquids consumed, you reduce the risk of developing electrolyte abnormalities such as low sodium. Please avoid any food or beverage product(s) which contain red or purple coloring.

MENU FOR CLEAR LIQUID DIET

<u>Breakfast</u>	<u>Lunch</u>	<u>Dinner</u>	
White Cranberry Juice	Chicken broth	Chicken broth	
Gelatin dessert	Apple Juice	White Grape Juice	
Tea/coffee (no milk)	Sprite, 7up, Ginger Ale	Gelatin dessert	
Gatorade or similar sports	Fruit-flavored ice	Sprite, 7up, Ginger Ale	
drink	Tea/coffee (no milk)	Tea/coffee (no milk)	

Instructions continue on the next page

TWO-DAY PREPARATION INSTRUCTIONS PRIOR TO COLONOSCOPY USING SuFLAVE

THREE DAYS PRIOR TO YOUR PROCEDURE: At 4:00 pm take pouch A and disposable container, add lukewarm water to the fill line of disposable container. Take pouch B and 2nd disposable container and add lukewarm water to the fill line of disposable container. Shake till mixed and put in refrigerator. DO NOT FREEZE.

FIRST DOSE:

STEP 1: Two days prior to your procedure at 6:00 pm take out of the refrigerator disposable container A and drink 8 ounces of the solution every 15 minutes until all the liquid in the container is finished.

STEP 2: Drink an additional 16 ounces of clear liquids after the solution is finished. (Refer to clear liquid diet menu)

THE DAY PRIOR TO YOUR PROCEDURE: At 4:00 pm take pouch A and disposable container, add lukewarm water to the fill line of disposable container. Shake till mixed and put in refrigerator. DO NOT FREEZE.

SECOND DOSE:

STEP 1: One day prior to your procedure, at 6:00 pm take out of the refrigerator disposable container B and drink 8 ounces of the solution every 15 minutes until all the liquid in the container is finished.

STEP 2: Drink an additional 16 ounces of clear liquids after the solution is finished. (Refer to clear liquid diet menu)

THIRD DOSE:

STEP 1: Starting 6 hours prior to your procedure time take out of the refrigerator disposable container A and drink 8 ounces of the solution every 15 minutes until all the liquid in the container is finished.

STEP 2: Drink an additional 16 ounces of clear liquids after the solution is finished. (Refer to clear liquid diet menu) PLEASE DO NOT CONSUME ANY LIQUIDS OR MEDICATIONS AFTER THE TIME THAT CORRESPONDS TO 4 HOURS BEFORE YOUR PROCEDURE TIME.

On average, your bowel movements should be clear (clear = yellow or white liquid without solid or granular material) 3 hours after you finish drinking the solution. However, delays in the onset of bowel movements and/or becoming clear can occur several hours after completing the solution. If you are not clear at 2 hours after taking the third dose on the day of your procedure, please follow the instructions written below.

If you are unable to complete and/or tolerate the preparation for colonoscopy, or your bowel movements are not clear after completing written instructions, please follow these instructions:

Purchase the following (no prescription necessary) and begin this preparation 1/2 hour after the last glass of solution taken.

One bottle of fleet enema (without mineral oil or any type of oil)

If your bowel movements are not clear, then proceed with Fleet Enema as follows:

Instructions continue to the next page.

- 1. Apply one Fleet Enema (without mineral oil, or any other type of oil) per rectum and wait 30 minutes.
- 2. If your bowel movements are still not clear, fill the same enema bottle with warm water from the faucet. Then administer a warm water enema every 30 minutes per rectum until you have bowel movements which consist of clear yellow or clear white liquid. Do NOT excede more than 4 enemas.

If you still encounter significant difficulties with your preparation, please contact our office at (703) 444-4799. If you are forwarded to voicemail, follow directions to contact the doctor on call.

RYAN P. CRENSHAW, M.D. 21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799 OPERATIVE REQUEST/CONSENT

1.	procedure(s) along with surg	gical assistants selected eeding, perforation, and	Crenshaw (the "Practitioner") to perform by him: colonoscopy, possible biopsy. Red need for surgery has been explained. The bons on (the "Patient"):	isk of drug allergy,			
	Please print your name:						
2.	2. It has been clearly explained to me that during the course of this operation some other conditions that have been expected may present themselves. I recognize, that if such conditions are discovered it will be necessar do more than that which was specified in paragraph #1 above. I therefore authorize and request that the ab named Practitioner and his surgical assistants perform such surgical procedures which in their best professi judgment will be effective in their attempt to heal and/or diagnose. This includes, but is not limited to, patho and radiology. I further authorize Anesthesiologist to administer whatever anesthesia they feel is indicated authorize the use of blood transfusion(s) when attending personnel feel such is required.						
3.	I fully understand that this opic is guaranteed.	ion, is accompanied by some degree of ris	k and that no cure				
4.	The nature of my (or the patient's) condition, the nature of the procedure(s) listed under paragraph #1 above, the risks involved and whatever other choices are available to me (or the patient), if any, have been explained to me by the Practitioner. I have been given the opportunity to ask any questions that I may have regarding that explanation and my questions have been answered satisfactorily.						
5.	I am aware of the "Cancellation Policy" and understand that I will be held responsible for a \$250.00 fee if notice is not provided at least <u>5 business days</u> in advance of scheduled date for procedure.						
.6.	If your procedure is cancelled due to non-compliance with both verbal and written instructions given (for example, not complying with the clear liquid diet the day prior to your procedure), you will be charged the cancellation fee.						
7. It is the patient's responsibility to contact his/her insurance provider to check coverage for the requested procedure(s), as well as obtaining the necessary referrals. It is also the patient's responsibility to notify our office immediately if your insurance provider changes, otherwise, the patient will be held responsible for any charges for the requested procedure(s).							
Sig	nature of Patient	Date	Signature of Witness	Date			
Sigi	nature of Parent/Guardian	Date					
Ιŀ		•	ATEMENT proposed procedure to the patient, and/or of this procedure, and any alternative.	Relative/guardian,			
	-	Signature of Physi	cian Date				

Page 5 of 5 Patient Copy: Two-day SuFLAVE

RYAN P. CRENSHAW, M.D. 21135 WHITFIELD PLACE, SUITE 102, STERLING, VA 20165 (703) 444-4799 OPERATIVE REQUEST/CONSENT

11	nave personally explained, in no the major –	n-technical terms, the properisks or consequences of the Signature of Physician	his procedure, and any	alternative.	elative/guardian, wo-day SuFLAVE		
		PHYSICIAN'S STATE	MENT				
Signature of Parent/Guardian		Date					
Sig	nature of Patient	Date	Signature of	Witness	Date		
7.	It is the patient's responsible procedure(s), as well as obtain immediately if your insurance the requested procedure(s).	ing the necessary referrals	s. It is also the patient's	responsibility to	notify our office		
.6.	. If your procedure is cancelled due to non-compliance with both verbal and written instructions given (for example, not complying with the clear liquid diet the day prior to your procedure), you will be charged the cancellation fee.						
5.	I am aware of the "Cancellation not provided at least 5 busines	•	·		00 fee if notice is		
4.	4. The nature of my (or the patient's) condition, the nature of the procedure(s) listed under paragraph #1 above, the risks involved and whatever other choices are available to me (or the patient), if any, have been explained to me by the Practitioner. I have been given the opportunity to ask any questions that I may have regarding that explanation and my questions have been answered satisfactorily.						
3.	I fully understand that this op is guaranteed.	eration, like any operation	, is accompanied by soi	me degree of risk a	and that no cure		
2.	been expected may present to do more than that which was named Practitioner and his su judgment will be effective in to and radiology. I further author	to me that during the course of this operation some other conditions that have not themselves. I recognize, that if such conditions are discovered it will be necessary to a specified in paragraph #1 above. I therefore authorize and request that the above-urgical assistants perform such surgical procedures which in their best professional their attempt to heal and/or diagnose. This includes, but is not limited to, pathology orize Anesthesiologist to administer whatever anesthesia they feel is indicated and ansfusion(s) when attending personnel feel such is required.					
	Please print your name:			_			
1.	I hereby request, consent of procedure(s) along with surgion sedation, aspiration, bleeding me there is a small possibility	cal assistants selected by hi , perforation, and need for	m: colonoscopy, possik surgery has been expla	ole biopsy, risk of o	drug allergy, over		